



The effect of sports specific circuit training and high intensity interval training on aerobic capacity in male basketball players: The case of Addis Ababa male basketball players

Solomon Teka

MSc in Coaching Basketball & Teaching PE, PhD (Ass. Prof.) in Exercise Physiology, Addis Ababa, Ethiopia Department of Sports Science, Addis Ababa University, Ababa, Ethiopia

Abstract

The demands and nature of basketball competition involve repeat high intensity efforts within a game, and during multi-game national and international tournaments. These demands require that the fitness and conditional attributes of basketball players are well developed to negate the limiting aspects of cumulative fatigue. Basketball is a game that is typically performed in an intermittent manner, and includes numerous bursts of explosive activity. Thus, implementation of training methods that best develop aerobic and anaerobic fitness is of significant importance to basketball coaches and players. The Purpose of the present study was conducted with ultimate objective of identify and evaluate the effect of Sports Specific Circuit Training and High Intensity Interval Training on aerobic capacity in the case of Addis Ababa male basketball players.

Methods: employed were quasi experimental study design. A total of 30 subjects of basketball players with age group 18 - 27 years of male basketball players, having agility score of <12.5 and V – sit and reach scores plus or above half inch from the point of zero.

Participants: 15 subjects in the Group “A” received Sports Specific Circuit Training and 15 subjects in Group “B” received High Intensity Interval Training.

Outcome measures: Aerobic Capacity ($VO_2\max$).

Results: Statistical analysis was done by using Paired “t” and independent “t” test showed significant improvement in improving aerobic capacity ($VO_2\max$).

Conclusion: It is concluded that there is statistically significant improvement in aerobic capacity High Intensity Interval Training when compared to Sports Specific Circuit Training in male basketball players.

Keywords: agility, sports specific circuit training, aerobic capacity $vo_2\max$, high intensity interval training, basketball players

Introduction

The game of basketball is characterized by numerous bursts of explosive activity, including jumping, turning, accelerating, and short distance sprinting. Consequently, the game is typically performed in an intermittent manner, and includes rapid and frequent changes in the movement patterns of the players. For example, McInnes *et al.* (1998) ^[49] found that the number of all activities during an entire game was 997 ± 183 , with a change in movement category every two seconds. In this report a mean total of 105 ± 52 high-intensity runs (mean duration 1.7 sec) was recorded, resulting in one high intensity run every 21 seconds. A similar activity pattern during basketball games was reported by Bishop and Wright (2006) ^[1] and by Ben Abdelkrim *et al.* (2007) ^[2] for under-19 year-old and for adult professional players, respectively. Basketball has gained worldwide popularity and fascinated players and spectators with its dynamic characteristics as a team sport. In this sport, players cover about 4500–5000m during a 40-min game with a variety of multidirectional movements such as running, dribbling, and shuffling at variable velocities and jumping (Narazaki *et al.* (2008) ^[4]). Besides, basketball is characterized by speed and repeated changes in the direction of activities and movement, especially since May 2000 when the rules were modified (Delextrat A, Cohen D. (2008) ^[6]).

Aerobic power is the ability to deliver oxygen to the tissues during prolonged bouts of exercise ($VO_2\max$). The ability to condition an athlete in aerobic training drills allows for play at higher-intensity levels and for prolonged periods of time. High-level aerobic training increases the anaerobic threshold, enhances lactic acid clearance, and delays the onset of fatigue. Muscle strength and power provide explosive energy both to the upper and lower extremities. These components are essential to maximize performance in rebounding, positioning, jumping, sprinting, and driving to the basket. Muscular endurance is vital to allow for repeated power throughout the course of a play or longer. Aerobic endurance training in conjunction with muscle strengthening provides peak performance for short and long energy demands (Steven J, Faoasm D. (2005) ^[8]). Physical capacity of athletes is an important element of success in sports achievements. It involves a huge number of different capacities, with aerobic capacity being its major component. Aerobic capacity denotes a general extent of metabolic processes occurring in the human organism, and stands for a larger portion of the total energetic capacity. Maximal oxygen uptake ($VO_2\max$) refers to the intensity of aerobic processes, and actually represents the capacity of the organism to utilize at a certain moment the maximum amount of oxygen $VO_2\max$ is considered to be the

most important component of endurance performance (Živanić S, Životić VM, Mijić R, *et al.*).

Physiological basis of physical capacity of an organism incorporates the functional capacity of the organism to increase the level of metabolic processes in keeping with the requirement of physical effort being exposed to the metabolic processes in this sense mean the transformation of chemical energy into mechanical one (Bowers RW, Fox EL.(1984) ^[10].

Morgan and Adamson at the University of Leeds first developed circuit training in the 1950's. It is a versatile training method as it can be adapted for many different situations, sections of the population and fitness requirements, and can be used at any time of the year. While the exercises are normally laid out in a circular pattern, the pattern can be varied for motivational purposes to that of a star, square, semi-circle, V-shape, line or zigzag (Morgan RE, Adamson GT. (1961) ^[11].

Circuit training is a method of fitness training that is designed to develop general, all-round physical and cardiovascular fitness (Scholich M. (1991). The term 'sport-specific training' implies that exercises should mimic as much as possible the actions of the body during participation in a given sport. Specificity should not, however, be over emphasized when selecting resistance exercises because it could lead to imbalances. Consequently, finding a balance between general and specific exercises would be appropriate in a circuit (Dick FW. 2002 and Williams M. 1993 ^[13, 14].

High-intensity interval training (HIT) is characterized by repeated sessions of relatively brief, intermittent exercise, often performed with an "all out" effort or at an intensity close to that which elicits peak oxygen uptake (i.e., 90% of VO₂ peak). High-intensity interval training (HIT) for several weeks improves markers of aerobic energy metabolism, such as maximal aerobic capacity and the maximal activities of mitochondrial enzymes. Regular endurance training improves performance during tasks that rely mainly on aerobic energy metabolism, in large part by increasing the body's ability to transport and utilize oxygen and by altering substrate metabolism in working skeletal muscle (Gibala MJ. (2007) ^[15]. Sports performance is the result of the interplay of various mental and physical factors. One of the main factors to improve and maximize athletic performance is the development of physical abilities. The capabilities of fitness are strength, speed, endurance and flexibility.

Agility is difficult players' components to describe and quantify. Qualities that define an agile basketball player include the ability to change direction with minimal loss of velocity, accelerate and decelerate within a minimal distance, and good proprioceptive foot movement for balance and coordination for weight transfer during evasive movements. Of the two main contributors to agility in the model proposed by Young *et al.* (2002) ^[3], the factors that underpin change of direction speed outweigh the perceptual and decision making factors required to anticipate, or create a change of direction. Although strength and power are components within the model that influence change of direction speed, these attributes have only a trivial correlation with speed or effective change of direction. These motor tasks are movements that make agility an imperative skill for basketball players.

Flexibility has been defined in many ways. It is used the term joint mobility indicates the degree of bending the normal range or scope of a joint or series of joints. Several sit-and-reach tests (SRs) are commonly used in health-related and physical fitness test batteries to evaluate the hamstring and lower back flexibility. Flexibility and agility are also key factors that permit the quick stops, starts, and turns these athletes must perform as well as sport-specific skills that are essential to successfully compete at a national and international level (Tumilty DM. (1993) ^[19].

A simpler method than monitoring heart rate is the Rating of Perceived Exertion (RPE). RPE is a subjective measure and asks the exerciser to think about how hard they feel their body is working against a standardized scale (Suchomel A. (2010) ^[20]. Many researchers concluded that sports specific circuit training is beneficial for improving aerobic capacity in basketball players and high intensity interval training improves aerobic capacity in soccer players. The effect of sports specific circuit training and high intensity interval training on aerobic capacity in male basketball players is currently unclear. The purpose of this study, therefore, was to identify and evaluate the benefits of sports specific circuit training and high intensity interval training on aerobic capacity in male basketball players.

Materials and Methods

Subjects

The purpose of the present study was conducted with ultimate objective of identify and evaluate the effect of Sports Specific Circuit Training and High Intensity Interval Training on aerobic capacity in the case of Addis Ababa male basketball players. This is a Quasi experimental, comparative type study done in Sports Authority of Addis Ababa Basketball Federation, Arat Kilo Educational Center (Arat Kilo Timihirt Maekel) Gymnasium, Riche Ethiopian Olympic Solidarity gymnasium and Ethiopian Basketball Federation Mini Stadium in Addis Ababa, Ethiopia. The players studied were basketball players and trained on a daily basis. 30 off season basketball players were conveniently selected and explained clearly about the procedure and written informed consent was taken and explained them the whole information obtained from them will be kept confidential.

Male subjects with age 18 - 27 years, with sit and reach test above ½ inch and T – Agility test score 10.5sec and below and who are able to fulfil the pre-screening test were included in this study. Subjects who had a history of fractures and trauma for past one year, who had a history of sprains and strains for past three months, history of Cardio-respiratory problems, participating in competitive sports, undergoing any other exercise training were excluded from the study.

The subjects were considered prior to the study for the details of equality, age, type of birth. Before providing the Questionnaire, they were explained in detail about the study and the questionnaire. Pre-test for the subjects was done with VAS and female sexual index questionnaire which was given to the subjects and explained and were asked to fill. After that the study population was divided in to two groups conveniently, Group "A" subjects were allotted, Group "B" subjects were allotted with the treatment of scar mobilization.

Procedure

Total of 30 Subjects were divided in to 2 groups by convenient sampling 15 in each group. Subjects were taken up for the study after they fulfilled the inclusion criteria. All subjects were evaluated prospectively in the ground and explained about sports specific circuit training to group “A” and High intensity interval training to group “B”. Written informed consent was obtained from them. All 2 groups were participated in the training program for 3 days a week 2 sessions per day for six weeks. Prior to the training the subjects VO₂max was measured by using 1Mile Run test Formula to predict VO₂max = 108.844 - 0.1636W - 1.438T - 0.1928H. (Where W = Weight in kg, T = Time for the one mile run and H = Heart Rate at the end of the run) and Treadmill VO₂max test VO₂max = 42 + (Time × 2) “Time” is the total time of the test expressed in minutes and fractions of a minute. Outcome measures were reassessed after 6th week of first assessment:

Group “A”

▪ **Sports specific circuit training**

The Group “A” underwent sports specific circuit training was carried out on outdoor courts for four times with four minute intervals for every subject. Training protocol average running time of one circuit was 36 second and the total distance covered during on lap was approximately 108m. The proportion of the circuit considered ‘offence’ activity where a basketball was dribbled was 42%, while 58% was considered ‘defensive’ activity without the ball. 72% of the movement forward sprinting, 20% side shuffling, 8% back pedalling. One field goal attempt, one rebound, three vertical jump, one pivot, and 14 changes of direction were completed during one repeat of the circuit.

Sports specific circuit training includes

1 - 2 Forward sprints; 2 Pivot left; 2-3 Shuffle left; 3-4 Shuffle right; 4-5. Shuffle left; 5-6 Shuffle right; 6-7 Run into vertical jump collect ball upon landing. 7-10 Speed dribble ball around the cones; 10-11 Speed dribbles; 11-12 Speed dribbles; 12-13 Layup attempt; 13 Layup rebound; 13-14 Speed dribbles; 14-15 Forward sprint; 15-1 back peddles circuit completes.

Group “B”

▪ **High intensity interval training protocol**

The Group “B” undergoes 6 weeks duration of High intensity interval training 3 days per week and 2 sessions per day on treadmill. The duration of training 64 minutes in this 32 minutes training in the morning and 32 min training in the evening, the intensity and speed based on rate of perceived exertion.

HIIT training session includes

Time intensity/speed perceived exertion 5min.; Warm up at an easy-moderate pace/4-5; 4min. gradually increase your speed, incline and/or resistance until you’re working at an intense pace. You should be out of your comfort zone and breathing hard, but able to sustain this level for the full 4 minutes/ 8-9; 2 min. Reduce speed to a very light pace to fully recover /3-4; 4min. Increase your speed, incline and/or resistance until you’re working at an intense pace/ 8-9; 2 min. Reduce speed to a very light pace to fully recover /3-4; 4 min. Increase your speed, incline and/or resistance until you’re working at an intense pace/ 8-9; 2 min. Reduce speed to a very light pace to fully recover /3-4; 4 min. Increase your speed, incline and/or resistance until you’re working at an intense pace/ 8-9; 5 min. Cool down at a very easy/3-4.

Results

From the 30 subjects analysed there is no dropout and all 30 subjects were taken with post-test assessment. The pre-test and post-test of 1 Mile run test and Treadmill VO₂max among Group “A” subjects were tabulated in Table 1. There was a statistical difference between pre-test and post-test 1 Mile run test in group “A” subjects trained with sports specific circuit training, p<0.05, mean of 1 Mile run test score before the sports specific circuit training was 57.8155 which was increased significantly to mean of 63.847 and there was a statistical difference between pre-test and post Treadmill VO₂max test in group “A” subjects trained with sports specific circuit training with p<0.05, mean of Treadmill VO₂max before the training was 59.003 which was increased significantly to mean of 62.962 after sports specific circuit training.

Table 1: Comparison of pre-test and post-test values of 1 Mile run test and Treadmill VO₂max in Sport Specific Circuit Training Group “A”.

Group “A”		Pre-Test		Post-Test		t-values	Significance
		Mean	SD	Mean	SD		
Pair 1	1 Mile run test	57.8155	2.778	63.847	2.940	-8.987	0.000
Pair 2	Treadmill VO ₂ max test	59.003	1.557	62.962	2.312	-14.374	0.000

Pre-test and post-test of 1 Mile run test and Treadmill VO₂max among Group “B” subjects were tabulated in Table 2. There was a statistical difference between pre-test and post 1 Mile run test in group “B” subjects treated with high intensity interval training p<0.05, mean of 1 Mile run test score before the high intensity interval training was 60.49 which was increased significantly to mean of 64.888 and there

was a statistical difference between pre-test and post Treadmill VO₂max test in group “A” subjects trained with high intensity interval training with p<0.05, mean of Treadmill VO₂max before the training was 60.822 which was increased significantly to mean of 65.017 after high intensity interval training.

Table 2: Comparison of pre-test and post-test values of 1 Mile run test and Treadmill VO₂Max in high intensity interval training Group “B”.

Group “B”		Pre-Test		Post-Test		t-values	Significance
		Mean	SD	Mean	SD		
Pair 3	1 Mile run test	60.49	2.03883	64.888	2.03903	-12.107	0.000
Pair 4	Treadmill VO ₂ max test	60.822	1.75662	65.017	2.09657	-19.407	0.000

Post-test values of 1 Mile run test and Treadmill VO₂max were compared between both the groups in Table 3, which shows the post-test mean value of 1 Mile run test in group “A” (63.8370) and in group “B” is (64.8885) which shows that there was no significant difference between both the

groups(p>0.05). This graph also shows the post-test mean value of Treadmill VO₂max in group “A” (62.9625) and in group “B” is (65.0170) which infers that there is a significant difference between both the groups(p<0.05).

Table 3: Comparison of post-test values of 1 Mile run test and Treadmill VO₂max Measure in between sport specific circuit training group “A” and high intensity interval training group “B”.

Group “B”		Pre-Test		Post-Test		t-values	Significance
		Mean	SD	Mean	SD		
1 Mile run test		63.8370	2.94082	64.8885	2.03903	-1.302	0.201
Treadmill VO ₂ max test		62.9625	2.31200	65.0170	2.09657	-2.944	0.006

Discussion

The game of basketball is recreational and a competitive game. It helps promotion of health, body control, alertness, co-ordination and team spirit. The major finding of this study is that specifically designed sport specific circuit training in basketball court and high intensity interval training on treadmill fulfils the criteria for aerobic capacity.

Results of this study showed that sports specific circuit training and High intensity interval training programs was more beneficial in terms of Aerobic capacity for the basketball players (p < 0.05) (Tables 1 and 2). Several underlying mechanisms for the effect of sports specific circuit training and High intensity interval training programs on Aerobic capacity for the basketball players have been proposed.

The normal aerobic capacity (VO₂max) for a male basketball players ranges from 45-65 (ml/kg/min).High intensity interval training improves the aerobic capacity due to the High intensity intermittent training program increases oxidative enzyme activity in muscle.

High intensity interval training was done on the treadmill thrice a week and two sessions per day which VO₂max is a good indicator of fitness level and may change with training. However, if training is already optimal then genetic factors may limit the VO₂ max obtained. This test must thus be interpreted in relation to your training and performance potential.

Sports specific circuit training design was based on a previous design Sperlich B, Zinner C, Heilemann I, *et al.* and adapted to mimic as closely as possible the movement patterns of basketball match play. The training is very easy to practice in the Basketball court which improves the aerobic capacity. The basketball specific endurance circuit induced greater improvements in aerobic fitness, suggesting it was somewhat effective in improving aerobic fitness during the competitive season.

High intensity interval training (HIT) is defined as recurring sessions of exercise bouts at an “all out” effort level (greater than 90% of HR max or VO₂max) separated by periods of recovery. Its primary use for many years has been with elite

athletes. Exercise duration and frequency may lead to variability in training adaptations. The length of the bout, number of times repeated in one session, time of day the bout is executed, recovery time between bouts, and recovery time between the sessions could also contribute to changes in metabolic, cardiovascular, and pulmonary adaptations.

Anaerobic glycolysis is a system in which glycogen is converted to ATP for immediate use, typically less than 2-3 minutes. Changes in glycogen, such as increased resting glycogen content and a reduced rate of glycogen utilization during matched-work exercise, were found with HIT. Increased resting glycogen content should result in a delay in the onset of fatigue. Subjects trained every day for two weeks in one study which led to a significantly lowered rate of glycogenolysis, increased creatine kinase, pyruvate kinase, and lactate dehydrogenase activity.

High intensity exercise bouts less than 10 seconds are considered mainly anaerobic but if a bout lasts longer than 10 seconds a greater demand for energy will occur and cause an increased involvement from the aerobic energy systems. If a 30 second bout were to be repeated three more times, by the fourth bout, the energy would primarily be derived from oxidative metabolism. This could possibly be attributed to an increased rate of oxygen transport and utilization.

Sprint-training bouts of 10 seconds with 50-second rest periods found an increase in the peak rate of the sarcoplasmic reticulum Ca²⁺ release following training. High intensity interval training has also shown improved mechanisms contributing to oxygen delivery. Sprint interval training, greater vasodilation of the blood vessels occurs during exercise which could be a mechanism for a reduction in resistance and a possible increased blood flow.

The increased dilation caused a decrease in vascular resistance which was greatest in the white region, the fast twitch skeletal muscle, of the trained rats consequently leading to a proposed overall increase in blood flow capacity. An increase in blood flow through a decreased vessel resistance allows for a greater oxygen delivery to the working muscles which increases the oxygen available for extraction which in turn may increase

VO₂max Dunham C. (2008) [54].

The improvement in the aerobic capacity after the sport specific circuit training because the duration of the study was six weeks three days per week two sessions per day which is more than the previous study. In mother study the training was done on the competitive session but this study was done on the off season basketball players.

Although there is improvement in Aerobic capacity (VO₂max) within the groups “A” and “B” (Tables 1 and 2) but in between the groups “A” & “B” Aerobic capacity (VO₂max) improvements are significant in treadmill VO₂max when compare to the 1mile run test after the spot specific circuit training and high intensity interval training (Table 3). Reasons for no improvements in Aerobic capacity (VO₂max) in 1 Mile Run test after trainings was the Players training programs were undergone on different surfaces and environment could be the initial, and the duration of the trainings were not similar at the baseline. For this reason there is no significant difference between sports specific circuit training and high intensity interval training in 1 mile run test values Aerobic capacity but in Tread mill VO₂max values shows improvement in aerobic capacity in Male Basketball Players (Table 3). Hence there is a significant difference in improving aerobic capacity after High intensity interval training when compared to Sports specific circuit training.

Conclusion

This study concludes that there is a significant difference in improving aerobic capacity after High intensity interval training when compared with Sports specific circuit training. As the tests result indicated that there was progressive improvement in performance from pretest to posttest on intervention groups of High intensity interval training and the circuit training.

A partially selective effect of the different kinds of training on physiological measures raises the possibility of prescribing training to correct weaknesses in these measures. On the basis of the existing research one can tentatively recommend adding or increasing explosive resistance training for a player with a poor economy and/or poor anaerobic capacity, and adding or increasing maximal intervals for a player with a poor VO₂max.

Acknowledgement

The authors thank the coaching staff and the players of the Addis Ababa Basketball clubs/Teams for their participation in the study and also wish to thank the Addis Ababa Basketball Federation organized the clubs for its cooperation in the study.

References

1. Bishop D, Write C. Time-motion analysis of professional basketball to determine the relationship between three activity profiles: high, medium and low intensity and the length of time spent on court. *Int J Per Analysis Sport*. 2006; 6:130-138.
2. Ben Abdelkerim N, El Fazza S, El Ati J. Time-motion analysis and physiological data of elite under 19 – year old basketball players during competition. *Br. J Sports Med*. 2007; 41:69-75.
3. Young, WB, James, R, and Montgomery, I. Is muscle power related to running speed with changes of direction? *The Journal of Sports Medicine and Physical Fitness*, 2002; 42(3):282-8.
4. Narazaki K, Berg K, Stergiou N, *et al*. Physiological demands of competitive basketball. *Scand. J Med. Sci. Sports*, 2008.
5. McInnes SE, Carlson JS, Jones CJ, *et al*. The physiological load imposed on basketball players during competition. *J Sports. Sci*. 1995; 13(5):387-397.
6. Delextrat A, Cohen D. Physiological testing of basketball players towards a standard evaluation of anaerobic fitness. *J Strength. Cond. Res*. 2008; 22(4):1066-1072.
7. Steingard PM. Basketball injuries. *Gin. Sports. Med*, 1993, 12.
8. Steven J, Faoasm D. Principles of manual sports medicine, Lippincott Williams & Wilkins, USA, 2005.
9. Živanić S, Životić VM, Mijić R *et al*. Aerobna sposobnost injena procena Astrandovim testom opterećenja na biciklrgometru. Beograd; Udruženje za medicinu sporta Srbijehhh.
10. Bowers RW, Fox EL. *Sports Physiology*, Mc- Graw-Hill, USA, 1984).
11. Morgan RE, Adamson GT. *Circuit Training*. Bell and Sons Ltd, London, 1961.
12. Scholich M. *Circuit training for all sports methodology of effective fitness training*, Sport Books, Toronto, USA, 1994.
13. Dick FW. *Sports training principles*, Human Kinetics Champaign, Illinois, USA, 2002.
14. Williams M. *Lifetime fitness and wellness*, Brown and Benchmark, Iona, 1993.
15. Gibala MJ. High Intensity Interval Training: New Insights. *Sports. Sci. Exch*. 2007; 20(2):1-6.
16. Bakirtzoglou panteleimon, ioannou panagiotis and bakirtzoglou fotis evaluation of hamstring flexibility by using two different measuring instruments.
17. Brown AE. The reliability and validity of the lane agility test for collegiate basketball players. *Hum. Perf*, 2012, 1-32.
18. Raven PB, Gettman LR, Pollock ML, *et al*. A physiological evaluation of professional soccer players. *Br. J Sports. Med*. 1976; 10(1):209-216.
19. Tumilty DM. Physiological characteristics of elite soccer players. *Sports. Med*. 1993; 16(1):80-96.
20. Suchomel A. Aerobic capacity and exercise intensity on different player levels in tennis. *Sport. Sci. Rev*. 2010; 19(1):1-2.
21. Liu J. Physiological responses of college male basketball athletes’ sport-specific aerobic interval training IEA 217. 2013; (1):691-697.
22. Marinković D, Pavlović S. The differences in aerobic capacity of basketball players in different playing positions *Series Phys. Edu. Sport*. 2013; 11(1):73-80.
23. Rad LS and Gholami M. Effects of changes in volume and intensity of exercise training on vo2 max in young females. *Ann. Biol. Res*. 2012; 3(4):1821-1827.
24. Karahan M. The effect of skill-based maximal intensity interval training on aerobic and anaerobic performance of female futsal players. *Biol. Sport*. 2012; 29(1):223-227.
25. Hottenrott K, Ludyga S, Schulze S. Effects of high

- intensity training and continuous endurance training on aerobic capacity and body composition in recreationally active runners. *J Sports. Sci. Med.* 2012; 11(1):483-488.
26. Ayala F, Baranda SD, Croix P, *et al.* Reliability and validity of sit-and-reach tests: systematic review. *Eur. J Appl. Physiol.* 2012; 5(1):57-66.
 27. Dunham C, Harms CA. Effects of high-intensity interval training on pulmonary function. *Eur. J Appl. Physiol.* 2012; 112(8):3061-3068.
 28. Carr N. The effect of high intensity interval training on vo2 peak and performance in trained high school rowers. *Int. J Sports. Physiol. Perform.* 2011; 4(1):110-121.
 29. Akilan N, Muthusubramanian J, Chittibabu B. Effect of sports specific circuit training on aerobic capacity of high school male basketball players during competitive season. *Asian. J Sci. Technol.* 2011; 1(12):073-075.
 30. Sporis G, Jukic I, Milanovic L, *et al.* Reliability and factorial validity of agility tests for soccer players. *J Strength. Cond. Res.* 2010; 24(3):679-686.
 31. Driller MW, Fell JW, Gregory JR, *et al.* The effects of high-intensity interval training in well-trained rowers. *Int. J Sports. Physiol. Perform.* 2009; 4(1):110-121.
 32. Narazaki K, Berg K, Stergiou N, *et al.* Physiological demands of competitive basketball. *Scand. J Med. Sci. Sports.* 2009; 19(3):425-432.
 33. Gormley SE, Swain DP, Gandrakota R. Effect of intensity of aerobic training on vo2max. *Med. Sci. Sports. Exerc.* 2008; 40(7):1336-1343.
 34. Helgerud J, Haydal K, Wang E, *et al.* Aerobic high-intensity intervals improve VO2max more than moderate training. *Med. Sci. Sports. Exerc.* 2007; 39(4):665-671.
 35. Stone N. Physiological response to sport specific interval training in high school male basketball players, School of recreation and sports, New Zealand, 2007.
 36. Chamari K, Hachana Y, Kaouech F *et al.* Endurance training and testing with the ball in young elite soccer players. *Br. J Sports. Med.* 2005; 39(1):24-28.
 37. Millan KM, Helgerud J, Donald RM, *et al.* Physiological adaptations to soccer specific endurance training in professional youth soccer players. *Br. J Sports. Med.* 2005; 39(1):273-277.
 38. Hoff J. Training and testing physical capacities for elite soccer players. *J Sport. Sci.* 2005; 23(6):573-582.
 39. Sallet P, Perrier D, Ferret JM, *et al.* Physiological differences in professional basketball players as a function of playing position and level of play. *J Sports. Med. Phys. Fitness.* 2005; 45(1):291-294.
 40. Dalleck LC, Kravitz L, Robergs RA. Maximal exercise testing using the elliptical cross-trainer and treadmill. *J Int. Elect.* 2004; 7(3):94-101.
 41. Hoff J, Wisløff U, Engen LC, *et al.* Soccer specific aerobic endurance training. *Br. J Sports Med.* 2002; 36(1):218-221.
 42. Finn C. Effects of high-intensity intermittent training on maximum oxygen uptake and endurance performance. *Sportscience.* 2001; 5(1):1-7.
 43. Kainoa P, Madole K, Garhammer J *et al.* Reliability and validity of the t-test as a measure of agility, leg power and leg speed in college-aged men and women. *J Strength. Cond. Res.* 2000; 14(4).
 44. Izumi T, Kouji N, Motoki K *et al.* Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO2max. *Med. Sci. Sports. Exerc.* 1996; 28(10):1327-1330.
 45. Dougall JDM, Wenger HA, Green HJ. Physiological testing of the high-performance athlete, Human Kinetics, USA, 1991.
 46. Dalleck LC, Kravitz L, Robergs RA. Maximal exercise testing using the elliptical cross-trainer and treadmill. *J Int. Elect.* 2004; 7(3):94-101.
 47. Chan D. Fitness testing assignment: Basketball the sporting supplement, 1999.
 48. Rodas G, Ventura JL, Cadefau JA *et al.* A short training programme for the rapid improvement of both aerobic and anaerobic metabolism. *Eur. J Appl. Physiol.* 2000; 2(5):480-486.
 49. McInnes SE, Maia J. game intensity in basketball. An interactionist view linking time – motion analysis, lactate concentration and heart rate *Coaching Sport Sci J.* 1998; 3:26-30.
 50. Garrick L. Health's Disease and Condition content.
 51. Sperlich B, Zinner C, Heilemann I, *et al.* High-intensity interval training improves VO2peak, maximal lactate accumulation, time trial and competition performance in 9–11-year-old swimmers. *Eur. J Appl. Physiol.* 2012; 110(5):1029-1036.
 52. Stellingwerff T. Explanation of Lactate Threshold and Vo2max.
 53. Smith M. Basketball skill test for the big men. 2004; 7(1), 59-60.
 54. Dunham C. The effects of high intensity interval training on pulmonary function. *Eur. J Appl. Physiology.* 2008; 12(8):3061-3068.