



Studies on complexation and mechanism of the interaction of metal complexes of amino acid with transition metals and determination of stability constant

Chitrendu Ji, Priyanka Sharma

Assistant Professor, Department of Chemistry, Future Engineering College, Bareilly, Uttar Pradesh, India

Abstract

It is notable that some metal particles are basic for living life forms. They assume vital parts in the combination and transport of natural atoms and in catalyzing corrosive base and redox procedures of organic frameworks. In metalloenzymes metal particles are normally bound by the side chain contributor elements of the peptides. The N(1)/N(3) contributor particles of histidyl and carboxylate gatherings of the aspartyl and glutamyl deposits are particularly favored restricting destinations of metalloenzymes.

Keywords: transition metal, schiff, amino acid

1. Introduction

The part of the side chain carboxylate bunches on the metal particle coordination of peptides and peptide subordinates has just been generally considered. The information demonstrates that the impact of the carboxylate bunches on the unpredictable development forms relies upon the number and area of these amino acids in the peptide chain. The impact of the area of these amino acids in the peptide chain on the mind boggling arrangement forms has just been generally considered. However the nearness of carboxylate bunches in the side chain changes the charge of the ligands and the buildings. We have explored the copper (II) and nickel (II) buildings of di-, tri- and tetra-peptides containing one, two or more aspartic or potentially glutamic acids. We have contemplated the impact of β - and γ -carboxylate amass and also the impact of expanded negative charge of buildings. The carboxylate assemble in the side chain of aspartic and glutamic corrosive likewise can be associated with amide official. The peptide-amide securities framed by the β - and γ -carboxylic gatherings of aspartic and glutamic acids are isolated from the peptide spine giving new coordination situations to metal particles. Lysine has comparative property; the ϵ -amino gathering can likewise shape a peptide bond with the carboxylic gathering of another amino corrosive, as on account of Asp- ϵ -Lys, or both amino gatherings of lysine can be associated with amide bound like on account of Gly-Lys (Gly). We have researched the copper (II), nickel (II) and zinc (II) buildings of these two peptides. We have examined the impact of the peptide bond shaped by ϵ -amino gathering. The investigations of ligands containing bis (imidazol-2-yl) assemble began around 15 years back. They examined the coordination science of the most straightforward ligands, at that point the N-secured tripeptide subsidiaries and the impact of the gatherings in chelatable position on the amino corrosive and peptide subordinates. Be that as it may, the impact of carboxylate assemble in the side chain were not known. We have explored the coordination science of ligands containing bis(imidazol-2-yl) aggregate on the C-end and a carboxylic

gathering in the side chain contrasting and the easiest dipeptides and the most straightforward amino corrosive subordinates of bis (imidazol-2-yl)methane. In addition these ligands can imitate the dynamic site of metalloenzymes. Consequently examines were proceeded into two bearings. As a matter of first importance we have researched the coordination science of bis (imidazol-2-yl) subordinates containing aspartic and glutamic corrosive. We have considered the impact of the side chain carboxylate amass on the complexation of amino corrosive bis (imidazol-2-yl)methyl subsidiaries through the ligands α -aspartyl-bis(imidazol-2-yl)methylamine (α -Asp-BIMA), α -glutamyl-bis (imidazol-2-yl)methylamine (α -Glu-BIMA) and γ -glutamyl-bis (imidazol-2-yl)methylamine (γ -Glu-BIMA). Then again we endeavored to copy Cu, Zn-SOD with different peptides.

2. Materials and Methods

In the dynamic site of Cu, Zn-SOD both restricting locales specified before are discovered; the particles are facilitated by the carboxylate gathering of aspartic corrosive and the imidazole nitrogens of histidine. The protein part of the compound which is incorporated into the metal particle restricting can be mirrored with oligopeptides. Hence we orchestrated peptides containing histidine and aspartic corrosive (Ac-His-Ala-His-Val-His-NH₂, Ac-His-Val-His-Ala-His-NH₂, Ac-His-ProHis-Ala-His-NH₂, Ac-His-Ala-His-Pro-His-NH₂, Ac-His-Val-Gly-Asp-His-NH₂ and Ac-HisHis-Val-Gly-Asp-NH₂). We portrayed the coordination science of these peptides with copper (II) particles.

3. Review of Literatures

Amino acids are of unique significance among the other concoction substances since they shape the essential constituents of living beings. It is basic to know the properties of amino acids with a specific end goal to comprehend and clarify their conduct and the blend of peptides, proteins and compounds in living life forms. It is realized that the

responses of peptides, proteins and catalysts with metal particles are of biochemical significance however they are yet to be completely comprehended. The clarification of these wonders in the natural frameworks can be conceivable just by the assurance of protonation constants of the amino acids and in addition their dependability constants, which are a measure of their propensity to shape edifices with other metal particles a medium like that of organic frameworks. It is watched that the vast majority of these conclusions have been done in fluid media (Norman and Doody, 1954; Norman *et al.*, 1958; Datta *et al.*, 1959; Singh and Srivastava, 1972; Gergely *et al.*, 1975; Robert *et al.*, 1985; Arbad and Sahaprdar, 1986) [1, 6, 4, 5, 7]. This might be credited to the across the board conviction that "in vivo" media are very much spoken to by the watery media. As of late, notwithstanding, it has been accounted for that watery media are not appropriate for organic in vivo responses, despite what might be expected the natural media indicate lipophilic character and are better spoken to by non-fluid media (Hughes *et al.*, 1986) [8]. The quantity of concentrates in non-watery media is extremely constrained in writing (Norman *et al.*, 1956; Rajashekar *et al.*, 1986) [2, 9]

4. Amino Acid Derivatives of bis(imidazol-2-yl)methyl Residue

The impact of side chain carboxylate aggregate on the complexation of amino corrosive bis(imidazol-2-yl) subsidiaries by means of the ligands α -Asp-BIMA, α -Glu-BIMA, γ -GluBIMA and for correlation β -Ala-BIMA.

With copper(II) particles the intricate development forms begin in exceptionally acidic arrangement ($\text{pH} \leq 2$), and the bis(imidazol-2-yl)methyl buildup is the selective restricting site in this pH area. The side chain contributor gatherings of amino acids frame six-or five membered chelate rings with terminal amino gathering and this outcomes in the arrangement of isomers of dinuclear edifices with ligand spanning. It was seen on account of α -Asp-BIMA (Scheme 5.a and 5.b) and γ -GluBIMA. For all amino corrosive subordinates, in which the terminal amino and the deprotonated amide nitrogen can shape five-or six membered chelates, the deprotonation of amide nitrogen happens at somewhat acidic pH (above pH 6), and the coordination of (NH_2, N^- , Im(N)) contributor set can break the bis(imidazol-2-yl)methyl coordination mode. It brings about the arrangement of dinuclear $\text{Cu}_2\text{L}_2\text{H}^- 2$ species, in which one of the imidazole rings of bis(imidazol-2-yl)methyl moiety serves a scaffold (Scheme 5.c). This procedure, be that as it may, does not happen on account of γ -Glu-BIMA, in which the amino and amide nitrogen could shape just seven membered chelate ring.

Another base expending procedure can be seen in soluble arrangement. This procedure can be clarified by blended hydroxo complex arrangement on account of amino corrosive subordinates with non-planning side chain (β -Ala-BIMA), and the deprotonation of pyrrole write nitrogen of ligands containing side chain giver gathering (α -Asp-BIMA and α -Glu-BIMA). As a result, the nearness of side chain giver gathering can keep the hydrolysis of copper (II) complex, presumably in view of its feeble collaboration with copper(II) particle. The deprotonation of the pyrrole write nitrogen makes another metal particle restricting site, which prompts

the presence of a trinuclear $\text{Cu}_3\text{L}_2\text{H}^- 4$ complex with contrarily charged imidazolate connect in the Cu (II)- α -Asp-BIMA (Scheme 5.d) and Cu(II)- α -Glu-BIMA frameworks at 3:2 molar proportion. The development of trinuclear species likewise can be assumed for Cu (II)- β -Ala-BIMA at abundance of metal particle. 3.3. The stoichiometries of the nickel (II) and zinc (II) buildings are fundamentally the same as those of copper(II) edifices, however these species have much lower thermodynamic secure qualities, which moves the perplexing development responses to higher pH esteems. As a result, the grouping of protonated buildings is much lower, particularly on account of zinc(II), and the security of the edifices diminishes in the request $\text{Cu(II)} > \text{Ni(II)} > \text{Zn(II)}$ as per the Irving-Williams arrangement.

Ccoordination of 4N giver molecules, while the customary octahedral geometry of nickel(II) and zinc(II) buildings gives a more adaptable coordination condition. In this way the measure of bis(ligand) buildings is higher. Then again, the moderately high steadiness of bis(ligand) edifices marginally smothers amide deprotonation in the relating nickel(II) and zinc(II) buildings when contrasted with that of copper(II) species. The convergence of dimer edifices with these particles are additionally lower, and in the Ni(II)- α -Asp-BIMA framework following the deprotonation of amide nitrogen a monomer qe dimer balance exists.

On account of α -Glu-BIMA and γ -Glu-BIMA other than deprotonation macro constants we have additionally figured deprotonation micro constants of these ligands. In α -Glu-BIMA the deprotonation grouping is Im1 d - COOH d Im2 d - NH_3^+ and the deprotonation forms fundamentally cover. In γ -Glu-BIMA this arrangement is COOH d Im1 d Im2 d - NH_3^+ and the deprotonation forms are particular.

5. Conclusion

The principle objective of this work was to uncover the collaboration between some change metal particles and amino corrosive and peptides subordinates. The contemplated ligands contain carboxylate and additionally imidazole bunches in their side chains, which are the principle restricting destinations of metalloenzymes

References

1. Norman CL, Doody Copper E. (II) and zinc complexes of some amino acids and glycyloglycine J Am. Chem. Soc. 1954; 76:221-225.
2. Norman CL, Norman JM, White RL. Yoesh some metal complexes of glycine and valine J Am. Chem. Soc. 1956; 78:5218-5222.
3. Singh Srivastava MK, Singh MN, *et al.* Srivastava Mixed ligand amino acid chelates of copper(II) and Ni(II) aspartates J. Inorg. Nucl. Chem. 1972; 35:2433-2440.
4. Gergely Gergely A, Nogypal L, Farkas E. *et al.* Thermodynamic relations of parent and mixed complexes of asparagines and glutamine with copper(II) J Inorg. Nucl. Chem. 1975; 37:551-555.
5. Robert Robert M, Motekaits RJ, Martell AE, *et al.* Prediction of stability constants II. Metal chelates of natural alkyl amino acids and their synthetic analogs. Inorg. Chim. Acta. 1985; 103:73-82.
6. Datta. Datta SP, Leberman R, Rabin BR, *et al.* The

- chelation of metal ions by dipeptides and related substances part II. The complexes of some sarcosyl and leucyl compounds with some divalent metal ions. *Trans. Faraday Soc.* 1959; 55:1982-1987.
7. Arbad Sahaprdar BR, Arbad DV. Sahaprdar Stability of Cu(II) glycy-l-asparagine amino acids ternary complexes and thermodynamic parameters of their formation *Ind. J Chem.* 1986; 25A:253-255.
 8. Hughes LD, Hughes JJ, Bergan JE. Grabowski Amino acid chemistry in dipolar aprotic solvent: Dissociation constants and ambident reactivity *J Org. Chem.* 1986; 51:2579-2585.
 9. Rajashekar NV, Rajashekar G, Ganpa RK, *et al.* Lomprakash Formation constants of binary and ternary chelates of Cu(II) with 2-(*N,N*-diethylaminomethyl) benzimidazole and amino acids *Ind. J Chem A*, 1986; 25:394-395.
 10. Maslawska Chruscinki J, Maslawska L, *et al.* Chruscinki Potentiometric studies on mixed ligand complexes of Cu(III) with DL-methionine, DL-ethionine and other amino acids *Polyhedron.* 1986; 5:1135-1139.
 11. Taylor CP. Taylor the Role of Gabapentin D. Chadwick (Ed.), *New Trends in Epilepsy Management*, Royal Society of Medicine Services Ltd., London. 1993, 13.
 12. Sing MJ, Field P, Ferris JC, Runker RJ, Oles RG, Williams GM. Woodruff the antiepileptic agent gabapentin (Neurontin) possesses anxiolytic – Like antinociceptive actions that are reversed by D-serine *Psychopharmacol. Berl.* 1996; 127:1.
 13. Backonja Glanzman M, Backonja RL. Glanzman Gabapentin dosing for neuropathic pain: evidence from randomized, placebo-controlled clinical trials *Clin. Ther.* 2003; 25:81.